

# ZYMAFLORE® F15

Yeast for fruity and round red wines

*Qualified for the elaboration of products for direct human consumption in the field of the regulated use in Oenology.  
In accordance with the International Oenological Codex.*

## SPECIFICATIONS AND OENOLOGICAL PROPERTIES

**ZYMAFLORE® F15** is for the production of **fruity, well-balanced** red wines with good mouthfeel (high **glycerol** production). It is suitable for the vinification of musts with potentially **high alcohol concentrations**, especially Merlot, Cabernet Sauvignon and Zinfandel.


### FERMENTATION CHARACTERISTICS:

- Alcohol tolerance: up to 16 % vol.
- Fermentation temperature tolerance: 20 - 32°C
- Medium nitrogen requirements
- Low production of volatile acidity and H<sub>2</sub>S

### AROMATIC AND ORGANOLEPTIC CHARACTERISTICS :

- High glycerol production
- Good varietal expression

## EXPERIMENTAL RESULTS



Production of glycerol by different strains of yeast on the same must.

## PHYSICAL CHARACTERISTICS

Dehydrated yeast (vacuum-packed)

Aspect.....granular

## STANDARD ANALYSIS

Humidity (%) .....	< 8 %	<i>Staphylococcus</i> UFC/g.....	None
Living cells SADY UFC/g .....	>2.10 <sup>10</sup>	<i>Salmonella</i> UFC/25 g .....	None
Lactic acid bacteria UFC/g .....	< 10 <sup>5</sup>	Moulds UFC/g .....	< 10 <sup>3</sup>
Acetic acid bacteria UFC/g .....	< 10 <sup>4</sup>	Lead .....	< 2 ppm
Wild yeast UFC/g .....	< 10 <sup>5</sup>	Arsenic .....	< 3 ppm
Coliforms UFC/g .....	< 10 <sup>2</sup>	Mercury .....	< 1 ppm
<i>E. coli</i> UFC/g .....	None	Cadmium .....	< 1 ppm

## PROTOCOL FOR USE

### ŒNOLOGICAL CONDITIONS

- Please refer to the Technical Booklet “*Good alcoholic fermentation management*” for complete information on yeast addition timing and techniques, the key points of fermentation.

### DOSAGE

- 15 - 30 g/hL (150 - 300 ppm)

In the case of prefermentative cold maceration (cold soaking), it is recommended to add yeast at 5 g/hL during tank filling, in order to dominate the indigenous flora, then to complete with 15 to 20 g/hL at the end of maceration, before increasing the must temperature.

### IMPLEMENTATION

- Carefully follow the yeast rehydration protocol indicated on the packet.
- Avoid temperature differences exceeding 10°C between the must and the yeast during inoculation. Total yeast preparation time must not exceed 45 minutes.
- In the case of potentially high alcohol degree potential and to minimise volatile acidity formation, use DYNASTART® / SUPERSTART® ROUGE in the yeast rehydration water.

### STORAGE

- Store in original sealed packages, in a cool dry place (off the floor) in an odour-free environment.
- Optimal date of use: 4 years.

### PACKAGING

500 g vacuum bag. 10 kg box.

